

Information on measures and related costs in relation to species included on the Union list - *Lagarosiphon major*

This note has been drafted by IUCN within the framework of the contract No 07.0202/2017/763436/SER/ENV.D2 “Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species”. The information and views set out in this note do not necessarily reflect the official opinion of the Commission, or IUCN. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor IUCN or any person acting on the Commission’s behalf, including any authors or contributors of the notes themselves, may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. This document shall be cited as:

Hussner, A. 2019. Information on measures and related costs in relation to the species included on the Union list - *Lagarosiphon major*. Technical note prepared by IUCN for the European Commission.

Date of completion: 04/07/2019

Comments which could support improvement of this document are welcome. Please send your comments by e-mail to ENV-IAS@ec.europa.eu.

Species (scientific name)	<i>Lagarosiphon major</i> (Ridley) Moss
Species (common name)	Curly waterweed
Author(s)	Andreas Hussner, Duisburg, Germany
Date Completed	04/07/2019
Reviewer	Jonathan R. Newman, Waterland Management Ltd, UK; Manuel A. Duenas, CEH, Wallingford, UK

Summary

Highlight of measures that provide the most cost-effective options to prevent the introduction, achieve early detection, rapidly eradicate and manage the species, including significant gaps in information or knowledge to identify cost-effective measures.

Lagarosiphon major is an evergreen submerged aquatic plant species, native to South America. The species has been introduced into Europe and is present in several European countries (Hussner, 2012), where it spreads exclusively via fragments, as no seed production is reported yet. *Lagarosiphon major* grows in stagnant and running waters, forming dense monospecific beds.

The unintentional introduction of aquatic species often occurs due to mislabelled or contaminated plant material in trade (Brunel, 2009; Champion *et al.*, 2010; Hussner *et al.*, 2014). As such, in order to prevent unintentional introductions of *L. major*, comprehensive controls of imported plant material should be implemented at places of entry and sites of sale of imported aquatic plants, so as to identify mislabelled or contaminated material. DNA barcoding tools can be used to support identification of *L. major* during inspections.

The spread of invasive alien aquatic plants into new water bodies often occurs through the transport of plant fragments attached to water sport equipment, such as boats and trailers (Johnstone *et al.*, 1985; Johnson *et al.*, 2001). This spread can be reduced by increasing public awareness of this problem via public campaigns and engagement activities and also through the implementation or improvement of biosecurity measures, which should also be incorporated into the actions recommended through public campaigns.

Early detection of new infestations can be achieved by intensive surveying (including through citizen science and professional scuba divers) or by using novel tools like eDNA. Early detection of small infestations increases the likelihood of successful control measures to eradicate the species and, for small infestations, benthic barriers, suction dredging and hand weeding are potential control methods (Caffrey *et al.*, 2010, 2011; de Winton *et al.*, 2013). Larger infestations can be managed by a combination of mechanical and/or chemical control followed by hand weeding (Caffrey *et al.*, 2010, 2011).

Prevention of intentional introductions and spread – measures for preventing the species being introduced intentionally. **This table is repeated for each of the prevention measures identified.** *If the species is listed as an invasive alien species of Union concern, this table is not needed, as the measure applies anyway.*

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p><i>As the species is listed as an invasive alien species of Union concern, the following measures will automatically apply, in accordance with Article 7 of the EU IAS Regulation 1143/2014:</i></p> <p><i>Invasive alien species of Union concern shall not be intentionally:</i></p> <ul style="list-style-type: none"> <i>(a) brought into the territory of the Union, including transit under customs supervision;</i> <i>(b) kept, including in contained holding;</i> <i>(c) bred, including in contained holding;</i> <i>(d) transported to, from or within the Union, except for the transportation of species to facilities in the context of eradication;</i> <i>(e) placed on the market;</i> <i>(f) used or exchanged;</i> <i>(g) permitted to reproduce, grown or cultivated, including in contained holding; or</i> <i>(h) released into the environment.</i> <p><i>Also note that, in accordance with Article 15(1) – As of 2 January 2016, Member States should have in place fully functioning structures to carry out the official controls necessary to prevent the intentional introduction into the Union of invasive alien species of Union concern. Those official controls shall apply to the categories of goods falling within the Combined Nomenclature codes to which a reference is made in the Union list, pursuant to Article 4(5).]</i></p> <p><i>Therefore measures for the prevention of intentional introductions do not need to be discussed further in this technical note.</i></p>
--	--

Prevention of un-intentional introductions and spread – measures for preventing the species being introduced un-intentionally (cf. Article 13 of the IAS Regulation). **This table is repeated for each of the prevention measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Control measures to check for mislabelled or contaminated plant material</p> <p>The major pathway of introduction of the species is intentional via aquatic plant trade (addressed in the table above). Nevertheless, the un-intentional introduction of aquatic species also occurs due to mislabelled or contaminated plant material in trade (Brunel, 2009; Champion <i>et al.</i>, 2010; Hussner <i>et al.</i>, 2014). As <i>L. major</i> is primarily clonal, imported aquatic plant material needs to be inspected for vegetative diaspores and particularly fragments of <i>L. major</i> (and any other plant species from the list of invasive alien species of Union concern). However, the degree of contamination of the species within the import of aquatic plants is not known, nor is the degree of mislabelling.</p> <p>For the contamination of plant material with diaspores and fragments, DNA barcoding tools can be used to support identification of <i>L. major</i> during inspections. In terms of mislabelled plant material, providing identification guidance to customs inspectors (e.g. see that produced by the GB NNSS¹), and importers and sellers will help increase detection.</p> <p>In addition to the un-intentional introduction via contaminated and mislabelled plants, the introduction into countries might also take place via propagules of <i>L. major</i> attached to e.g. boats and trailers (see section on secondary spread for more detailed information). <i>Lagarosiphon major</i> has a high resistance to desiccation (Heidbüchel <i>et al.</i>, 2019) and may survive even long-term overland transport.</p>									
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>The measures to control imported plant material, in order to identify mislabelled or contaminated material, should be undertaken across the EU and installed at sites of entry (i.e. airports, ports), but also at sites of sale of imported aquatic plants (i.e. aquarium and plant shops), to identify any contamination of plants in trade with propagules of <i>L. major</i>.</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>		<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>		<p><i>Unknown</i></p>	<p>X</p>	
<p><i>Rationale:</i></p>										

¹ <http://www.nonnativespecies.org/factsheet/factsheet.cfm?speciesId=1888>

<p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>The effectiveness of such control measures depends on the effort and resources applied. However, the success of import controls is hard to quantify, as no information about the level of contamination of imported plant material with the target species is available, and the number of mislabelled imported plant material is unknown. Furthermore, the success of these control measures depends on various parameters, such as knowledge of the species by inspectors and other responsible authorities. A border control programme for aquatic weeds has been developed for New Zealand (Champion & Clayton, 2001), but no information about the success of this border control is available.</p>																											
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>Control sites must be installed for the long-term to prevent any unintentional introduction of <i>L. major</i>.</p>																											
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>While inspection capacity already exists across the EU, additional control efforts usually entail additional costs; the costs per species will, however, be significantly reduced, if control measures are implemented for all aquatic plant species of Union concern at the same time. The identification of a species from plant material and particularly from diaspores requires an excellent species knowledge of inspectors. DNA barcoding tools, which have been developed for other invasive alien aquatic plant (IAAPs) species of union concern, like <i>Hydrocotyle ranunculoides</i> (Van de Wiel <i>et al.</i>, 2009), are a valuable tool to improve the reliability of the determination of diaspores and plants, but have not been developed for <i>L. major</i> yet. Resources are required for and should be invested into this.</p>																											
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 898 1888 999"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> </table> <p><i>Rationale:</i> The implementation of import control measures requires a good species knowledge and/or valuable tools (e.g. DNA barcoding), and will help to limit the import of other invasive alien aquatic plant species into the EU, especially of relatives of <i>L. major</i> from the family of Hydrocharitaceae.</p> <p>Any restrictions of plant imports increase the costs for the import of plants, and can thus reduce the number of plants sold. However, the sale of native species can be supported and similar native plants can be sold as alternatives, as has been successfully initiated in Belgium as part of an agreement between representatives from the ornamental sector, public authorities and the scientific community (Halford <i>et al.</i>, 2014).</p>							<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X
<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>																							
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																						

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="638 150 1883 220"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Unacceptable</i></td> <td><input type="checkbox"/></td> </tr> </table> <p><i>Rationale:</i> The establishment of voluntary agreements between the ornamental sector and authorities in Belgium (Halford <i>et al.</i>, 2014) indicates that the awareness about the problem of invasive aquatic plants is high, and thus control measures should be accepted by traders. Nevertheless, increased costs for the traders (due to restrictions of import) will most likely reduce the acceptability of any prevention measure.</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>	
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>			
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>No data about the costs of implementation and maintenance of import controls are available for aquatic plants. Consequently, no information can be given about the cost-effectiveness of this measure. However, the intensive control of imported plant material will incur additional costs but, as mentioned above, it can be implemented as a general control of introduced plant material on plants prohibited from introduction and sale in accordance to EU Regulation 1143/2014.</p> <p>The costs of inaction are hard to quantify, though as <i>L. major</i> is able to grow in a wide range of freshwater habitats throughout the EU, the costs of inaction (and subsequent eradication or control) will likely be higher than the costs of implementing these prevention measures (Hussner <i>et al.</i>, 2017).</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="638 858 1924 928"> <tr> <td><i>Inconclusive</i></td> <td><input type="checkbox"/></td> <td><i>Unresolved</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Established but incomplete</i></td> <td><input type="checkbox"/></td> <td><i>Well established</i></td> <td><input type="checkbox"/></td> </tr> </table> <p><i>Rationale:</i> Comprehensive import control must be considered as a valuable tool to stop the unintentional introduction of <i>L. major</i> and other invasive aquatic plant species. There is a consensus that the prevention of introduction of invasive alien aquatic plant species is cheaper than their later management and control (Hussner <i>et al.</i>, 2017). However, and although identification tools for whole plants, as well as for any part of a plant (e.g. DNA barcoding) exist for aquatic plants on the species level, no information about the implementation, ongoing costs, success and failures of these tools exist.</p>	<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input checked="" type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input type="checkbox"/>
<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input checked="" type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input type="checkbox"/>		

Prevention of secondary spread of the species – measures for preventing the species spreading once they have been introduced (cf. Article 13 of the IAS Regulation). This table is repeated for each of the prevention measures identified.

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Public awareness campaigns for, and coupled with, improved biosecurity practices</p> <p>It has been documented that the spread of invasive alien aquatic plants into new water bodies largely depends on the transport of plant fragments attached to water sport equipment, transported from one water body to another (Johnstone <i>et al.</i>, 1985; Johnson <i>et al.</i>, 2001). Aquatic plants produce fragments either by allofragmentation (fragmentation caused by disturbances) or autofragmentation (self-induced fragmentation). The likelihood of successful spread increases with increasing number of fragments produced and transported into a new water body (Lockwood <i>et al.</i>, 2005).</p> <p>In general, the number of fragments which can be found in a water body is species specific (Heidbüchel <i>et al.</i>, 2019). The number of fragments produced by <i>L. major</i> is relatively low, and was found to be much lower than in <i>Elodea canadensis</i> and <i>Egeria densa</i> (Redekop <i>et al.</i>, 2016). Moreover, the likelihood for regeneration of fragments is affected by their type, as fragments with apical tip show significantly higher regeneration rates than fragments without apical tip (Heidbüchel & Hussner, 2019).</p> <p>The human-mediated un-intentional spread of <i>L. major</i> can occur via plant fragments attached to any water sport equipment, but boats and trailers are considered the major transport vectors for such fragments (Johnstone <i>et al.</i>, 1985). The survival time of the attached fragments depends on both weather conditions (the warmer and less humid, the shorter the survival period) and the clumping of fragments (single fragments have a shorter survival period than clumped patches of plant fragments). <i>Lagarosiphon major</i> fragments with apical tip keep viable even after >60 % water loss. Chlorophyll fluorescence was documented as a valuable tool for the determination of the viability of the fragments (Heidbüchel <i>et al.</i>, 2019).</p> <p>To stop the spread of invasive alien aquatic plant species by fragments attached to boats and trailers, various preventive methods are reported. Overland dispersal of fragments via watersport equipment can be reduced by increasing public awareness of this problem via public campaigns like “Stop Aquatic Hitchhikers” initiated in the USA², Canada³ or New Zealand (author’s observations), and the UK’s “Check Clean Dry”⁴ and “Be Plant Wise”⁵ campaigns. In general, the transport of viable fragments can also be reduced by undertaking several biosecurity measures (see below; Johnstone <i>et al.</i>, 1985; Barnes <i>et al.</i>, 2013; Anderson <i>et al.</i>, 2015; Cuthbert <i>et al.</i>, 2018;</p>
--	---

² https://www.fws.gov/fisheries/ans/pdf_files/Stop_Aquatic_Hitchhikers_factsheet.pdf

³ http://www.env.gov.bc.ca/fw/fish/regulations/docs/1011/fa_AquaticHitchhikers.pdf

⁴ www.nonnativespecies.org/checkcleandry/

⁵ www.nonnativespecies.org/beplantwise/

	<p>Hippolite & Kurapa, 2018; Crane <i>et al.</i>, 2019), which should be incorporated into the actions recommended through any public campaign:</p> <ul style="list-style-type: none"> (i) the creation of weed free haul-out areas to reduce the likelihood of fragments attaching to boats and trailers, (ii) the collection of all visible plant fragments from boats and trailers, <p>or killing the fragments by:</p> <ul style="list-style-type: none"> (iii) storing of boats and trailers prior to the release into new water bodies, (iv) placing the boat into a wash station, a heated water system or exposing fragments to steam, and (v) using aquatic disinfectants prior to the release of the trailered boats into a new water body. 							
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Public engagement activities are usually undertaken at a national level. In New Zealand, a public awareness campaign was brought to the public by informing various business organisations (e.g. outdoor, boat and fishing gear retailers and tourist attractions). Additionally, people were informed during local water related events, and even school classes were visited and both teachers and children were informed (Hippolite & Kurapa, 2018).</p> <p>For biosecurity strategies, weed free haul-out areas are in use in several waters in e.g. the Bay of Plenty region, New Zealand (author's observation). Moreover, in this region, portable boat wash stations were tested for the in-field use at boat ramps (Hippolite & Kurapa, 2018). As alternatives to washing stations, heated water tanks (Anderson <i>et al.</i>, 2015), steam exposure (Crane <i>et al.</i>, 2019) or aquatic disinfectant (Cuthbert <i>et al.</i>, 2018) have been successfully tested in a laboratory experiment to kill fragments, but no information on field trials is available.</p>							
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 874 1794 943"> <tr> <td data-bbox="645 874 927 943">Effectiveness of measures</td> <td data-bbox="927 874 1095 943"><i>Effective</i></td> <td data-bbox="1095 874 1153 943" style="text-align: center;">X</td> <td data-bbox="1153 874 1279 943"><i>Neutral</i></td> <td data-bbox="1279 874 1509 943"></td> <td data-bbox="1509 874 1736 943"><i>Ineffective</i></td> <td data-bbox="1736 874 1794 943"><i>Unknown</i></td> </tr> </table> <p><i>Rationale:</i></p> <p>Public campaigns have been initiated in several countries, but the efficiency of such measures is difficult to quantify. However, in the UK, the 'Check Clean Dry' campaign increased the numbers of general public following the recommended biosecurity procedures by 9% (and 14% in high risk user compliance; Burchnall, 2013). Moreover, a 40% reduction of biosecurity hazards were reported for anglers and canoeists who have heard of the 'Check Clean Dry' campaigns, compared to anglers and canoeists who have not heard about it (Anderson <i>et al.</i>, 2014).</p> <p>In the Bay of Plenty Region in New Zealand, 75-86% of the users have freshwater weeds knowledge and were able to name any pest species, and 74-82 % follow biosecurity campaign measures when moving their boat from one freshwater to another (Hippolite & Kurapa, 2018). Nevertheless, the efficiency of public campaigns will most likely depend on the way and intensity by which the public has been informed. A combination of on-site information posters and press releases are recommended.</p>	Effectiveness of measures	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>	<i>Unknown</i>
Effectiveness of measures	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>	<i>Unknown</i>		

	<p>The success of direct biosecurity measures will depend on various parameters, e.g. the resistance of plants and seeds to desiccation and heating (Barnes <i>et al.</i>, 2013; Anderson <i>et al.</i>, 2015). However, considering the strong evidence for the importance of human-mediated overland dispersal of plant fragments via water sports equipment (Johnstone <i>et al.</i>, 1985), measures to stop this vector of unintended spread are considered to have a high impact. For <i>L. major</i>, some of these measures seem to be quite effective (see examples below).</p>																					
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>For the sustainable success of public campaigns, they generally need to be applied in the long-term and must target all kinds of water users.</p> <p>For biosecurity control, the exposure of <i>L. major</i> fragments to 45°C for 1 hour resulted in 100% mortality. The time required for drying to kill fragments depends on weather conditions and clumping of fragments (Heidbüchel <i>et al.</i>, 2019). Exposure to the aquatic disinfectant Virasure for 2 min at 1% concentration or 1 min at 4% showed optimised degradation of <i>L. major</i> fragments (Cuthbert <i>et al.</i>, 2018). Direct steam exposure, at a distance of 2-3 cm to the fragments, results in the die off of fragments (Crane <i>et al.</i>, 2019).</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The costs of generating a public awareness campaign are relatively low compared to the costs of managing established IAAPs. Even though public awareness campaigns were initiated in a number of countries, no data about the required resources are available.</p> <p>The installation of net cages in lakes to create weed free areas requires scuba diving activity and ongoing maintenance, and fragments caught by the net cages must be removed and disposed of in an appropriate manner. For the installation of boat washing stations, heated water systems or a system for steam exposure, specific equipment and experienced workers are needed. Similarly, the use of aquatic disinfectant requires experienced workers.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1" data-bbox="645 1054 1888 1157"> <tr> <td>Environmental effects</td> <td>Positive</td> <td>X</td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td></td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> </table> <p><i>Rationale:</i> The described methods provide a general barrier for the unwanted spread of invasive alien aquatic plants in general, which has positive environmental effects. Biosecurity measures can potentially impact the spread of native organisms, but such negative impacts on native plants have not been reported yet. Although public campaigns generally have a positive social effect of raising awareness about other aquatic alien species, biosecurity measures affect the recreational human use of water bodies, creating a mixed social effect.</p>	Environmental effects	Positive	X	Neutral or mixed		Negative		Social effects	Positive		Neutral or mixed	X	Negative		Economic effects	Positive		Neutral or mixed		Negative	X
Environmental effects	Positive	X	Neutral or mixed		Negative																	
Social effects	Positive		Neutral or mixed	X	Negative																	
Economic effects	Positive		Neutral or mixed		Negative	X																

<p>rationale, with supporting evidence and examples if possible.</p>	<p>Biosecurity measures will also represent extra time and costs for boat owners.</p>															
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 233 1888 296"> <tr> <td data-bbox="640 233 925 296">Acceptability to stakeholders</td> <td data-bbox="925 233 1151 296"><i>Acceptable</i></td> <td data-bbox="1151 233 1211 296"></td> <td data-bbox="1211 233 1509 296"><i>Neutral or mixed</i></td> <td data-bbox="1509 233 1570 296">X</td> <td data-bbox="1570 233 1832 296"><i>Unacceptable</i></td> <td data-bbox="1832 233 1888 296"></td> </tr> </table> <p><i>Rationale:</i> While public campaigns are usually well accepted by stakeholders, the recommended active biosecurity measures (collecting fragments, washing, heating and disinfecting) will incur additional costs and take time, which might have a negative public perception, although this has not been investigated yet. The public campaign in itself can help to improve the understanding and acceptance of these measures.</p>								Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>		
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>											
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>In general, no data about the costs of such public campaigns are available, but the costs will vary largely according to the number of in field information (e.g. information boards) needed. The costs for the measures to prevent species spread by killing the fragments depend on the number of boats that are transported overland and the number of lakes infested with the target species.</p> <p>The costs of inaction are hard to quantify, but as <i>L. major</i> is able to grow in a wide range of freshwater habitats, the costs of inaction (and subsequent eradication or control) will be higher than the costs of implementing these measures (Hussner <i>et al.</i>, 2017).</p>															
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="640 943 1924 1007"> <tr> <td data-bbox="640 943 913 1007"><i>Inconclusive</i></td> <td data-bbox="913 943 965 1007"></td> <td data-bbox="965 943 1167 1007"><i>Unresolved</i></td> <td data-bbox="1167 943 1227 1007"></td> <td data-bbox="1227 943 1509 1007"><i>Established but incomplete</i></td> <td data-bbox="1509 943 1570 1007">X</td> <td data-bbox="1570 943 1868 1007"><i>Well established</i></td> <td data-bbox="1868 943 1924 1007"></td> </tr> </table> <p><i>Rationale:</i> Public campaigns to stop the unintentional, human mediated spread of <i>L. major</i> were implemented in e.g. New Zealand or the USA with good success (Hippolite & Kurapa, 2018). Moreover, for invasive submerged aquatic plant species which predominantly spread by fragments, like <i>L. major</i>, prevention of spread by implementing biosecurity measures to remove or kill fragments attached to boats and trailers is of high relevance. Boat washing stations have been, for example, successfully tested in the Bay of Plenty, New Zealand (Hippolite & Kurapa, 2018). In contrast, the use of e.g. aquatic disinfectants or hot steam exposure has only been tested under laboratory conditions (Cuthbert <i>et al.</i>, 2018; Crane <i>et al.</i>, 2019).</p>								<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>										

Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the early detection measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Surveying through citizen science</p> <p>The detection of early infestations of invasive alien aquatic plant species is crucial for the later likelihood for rapid eradication of the target species (Genovesi <i>et al.</i>, 2010; Hussner <i>et al.</i>, 2017). If newly introduced invasive alien aquatic plant species are detected early, the eradication of these new populations is achievable prior to their establishment, which drastically reduces the costs of eradication measures (Hussner <i>et al.</i>, 2017).</p> <p>Citizen science programmes are a useful tool for surveying new incursions of invasive aquatic plants. They can show a high accuracy of data collected (80–95% accuracy; Delaney <i>et al.</i>, 2008) but, in the case of submerged aquatic weeds, specialised citizen scientists, i.e. recreational scuba divers with experience in macrophyte mapping, are usually required to undertake records. In lakes, scuba divers are required to identify early infestations of submerged plants, while in small to medium sized rivers submerged plants can usually be identified from the shore.</p> <p>Citizen science projects can be led by citizen scientists or, most commonly, by scientists in participation initiatives. For example, in Germany, the Federal Agency for Nature Conservation initiated a joint project with recreational divers, informing about invasive alien aquatic plant species and providing information brochures and identification keys (http://www.neobiota.info/Neophyten.php). Such engagement activities could also reach out to other recreational user groups.</p> <p>A surveillance system to support early detection should also incorporate active monitoring of the species; see tables below on use of scuba divers and environmental DNA.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Citizen science initiatives with recreational divers, and other key groups, would be undertaken at a national scale and drastically increase the number of water bodies that could be investigated, in relation to more traditional surveying methods.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 151 1888 220"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td><i>X</i></td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td></td> <td><i>Unknown</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> The efficiency of citizen science projects depends on the species knowledge of the involved citizen scientists. In a citizen science project dealing with the distribution and abundance of crabs in intertidal zones, species identification reached an accuracy between 80 and 95% (Delaney <i>et al.</i>, 2008). However, there is often concern about the quality of data provided by citizen scientists (Hochachka <i>et al.</i>, 2012) and the high phenotypic plasticity of submerged plants like <i>L. major</i> makes species identification difficult. Therefore, active engagement with and provision of identification materials to key groups (e.g. recreational divers) will be critical to guarantee or improve detection accuracy.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	<i>X</i>	<i>Neutral</i>		<i>Ineffective</i>		<i>Unknown</i>													
<i>Effectiveness of measures</i>	<i>Effective</i>	<i>X</i>	<i>Neutral</i>		<i>Ineffective</i>		<i>Unknown</i>															
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>Early detection of invasive submerged aquatic plants requires comprehensive, repeated surveillance of the macrophyte communities.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The development of a comprehensive and accurate citizen science project requires a substantial coordination by a government or scientific body. The annual costs for running a citizen science project are approximately between 80,000 – 170,000 EUR (Roy <i>et al.</i>, 2012 in Newman & Duenas, 2017).</p> <p>Detailed species identification sheets and keys must be produced and provided to allow citizen scientists to identify the target species (e.g. Adriaens <i>et al.</i>, 2015). Data recording apps (incorporating verification of records) are also needed to enable citizen scientists to report geolocated data records of new infestations of the target species.</p> <p>Citizen science data recording apps already exist for reporting occurrences of invasive alien species of Union concern at the EU level (the JRC EASIN Invasive Alien Species Europe smartphone App⁶) and within some EU Member States (e.g. see GB NNESS smartphone apps⁷).</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p>	<table border="1" data-bbox="640 1107 1888 1214"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td><i>X</i></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td><i>X</i></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td><i>X</i></td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i></p>	<i>Environmental effects</i>	<i>Positive</i>	<i>X</i>	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>	<i>X</i>	<i>Neutral or mixed</i>		<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Negative</i>																	

⁶ <https://easin.jrc.ec.europa.eu/easin/CitizenScience/About>

⁷ <http://www.nonnativespecies.org/index.cfm?sectionid=81>

<p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>In general, citizen science projects cause greater awareness of existing and future environmental problems, and increase the likelihood of additional IAS being reported.</p>								
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Unacceptable</p>			
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>It is widely accepted that, in general, the costs of inaction will be much higher than those of early detection measures, as the control and eradication of large infestations of IAAPs is much more time consuming and costly (Hussner <i>et al.</i>, 2017). The cost-effectiveness of early detection measures has not been studied in detail yet, but will differ between species and habitats.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p>Inconclusive</p>		<p>Unresolved</p>		<p>Established but incomplete</p>	<p>X</p>	<p>Well established</p>		
	<p><i>Rationale:</i> Citizen science projects have been reported to provide accurate observations of different species; however, its application to <i>L. major</i> and other submerged aquatic plants that require expertise to be confidently identified is not yet established.</p>								

Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the early detection measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Surveying through eDNA</p> <p>A new valuable tool for early detection of organisms is the use of environmental DNA (eDNA) markers, allowing the detection of a target species in water bodies, when the eDNA concentration in the water reaches a detection threshold. The detection of a given species requires a species specific primer and, After a positive eDNA record, an active survey programme is needed to locate the target species within the water body. Until yet, no eDNA studies dealing with <i>L. major</i> are available, and probably no primer has been developed for this species, but this method has been successfully tested with other Hydrocharitaceae species (Matsuhashi <i>et al.</i>, 2016).</p>									
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>The use of eDNA allows for the fast analysis of waters from numerous water samples, drastically increasing the potential number of water bodies which can be investigated. For example, the detection of rare fish species, even in large >100 ha lakes, can be done within one day (Hussner <i>et al.</i>, unpublished).</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 826 1888 890"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td>X</td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td></td> <td><i>Unknown</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> The use of eDNA is a relatively new tool and only few studies have analysed the efficiency of this method for identification of invasive aquatic plants (Scriver <i>et al.</i>, 2015; Matsuhashi <i>et al.</i>, 2016). In general, eDNA analysis requires the development of species specific primers (Scriver <i>et al.</i>, 2015). In a recent study (Scriver <i>et al.</i>, 2015), ten aquatic plant species were successfully detected from water samples. Matsuhashi <i>et al.</i> (2016) tested the sensitivity of the eDNA method in the field and in an aquarium experiment using two Hydrocharitaceae, <i>Hydrilla verticillata</i> and <i>Egeria densa</i>. The authors document that both plants do not release constant amounts of eDNA, which makes the estimation of biomass difficult, but the method was found to be a valuable tool for the identification of the plant species within the water body. More studies are needed to evaluate the minimum biomass of submerged plants needed to get a high accuracy of the species detection.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>		<i>Unknown</i>	
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>		<i>Unknown</i>			
<p>Effort required</p>	<p>In general, eDNA samples can be taken whole year round, but no information exists on if sampling time affects the accuracy of the analysis for evergreen aquatic plant species. It is important to note that eDNA analysis provides only</p>									

<p>e.g. period of time over which measure needs to be applied to have results</p>	<p>presence/absence data for the water body tested, the records are not geolocated, and thus, after any positive result, additional macrophyte surveying is required to locate the species.</p>																											
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>eDNA studies require experienced scientists for both sampling and analysis. The costs of the analysis vary according to the number of different species which need to be identified from the sample, while the costs of sampling are the same. The sampling and analysing of water samples to identify the fish community in certain lakes cost ca. 2,000-3,000 EUR per lake (Hussner, unpublished).</p>																											
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>							
<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>																							
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
<p><i>Rationale:</i> Overall, during eDNA studies other IAAPs can be identified, which reduces the total cost of monitoring per species.</p>																												
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																					
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																							
<p><i>Rationale:</i> This measure should be acceptable to all stakeholder groups.</p>																												
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects 	<p>There is no information available.</p>																											

Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).								
Level of confidence on the information provided ²	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	<i>X</i>	<i>Well established</i>	
Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure	<p><i>Rationale:</i> eDNA is a new powerful tool for species presence and absence detection, but more studies are needed to evaluate the minimum biomass of submerged plants needed to get a high accuracy of the species detection.</p>							

Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the early detection measures identified.	
Measure description Provide a description of the measure, and identify its objective	<p>Active surveillance using scuba divers</p> <p>Particularly for early detection of small infestations of submerged weeds like <i>L. major</i> in deeper waters, or for the confirmation of new occurrence records (e.g. via citizen science or eDNA, see surveillance tables above), scuba diving seems to be the most appropriate method for in field identification. Within the EU, macrophyte surveying by scuba divers is widely used as part of regular monitoring programmes, in accordance to the Water Framework Directive (WFD), and could be used as part of an early detection programme.</p> <p>The use of cable connected sub cameras allows a more rapid survey of macrophyte communities in a larger scale, but is less sensitive than surveys by scuba divers (Van de Weyer <i>et al.</i>, 2007).</p>
Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please	<p>Using scuba diving as an active survey measure is time consuming and allows detailed mapping of only small areas. Moreover, even zero-detection surveys do not definitely mean that the investigated water body is free of the target species, as finding of small infestations by scuba diving is generally difficult, particularly in turbid waters (Anderson, 2005). The macrophyte mapping in accordance to the WFD is usually limited to small areas (usually transects) in large lakes >50 ha.</p>

provide examples, with areas (km ² or ha) if possible.																								
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Effectiveness of measure</i></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	<i>Effectiveness of measure</i>									<table border="1"> <tr> <td><i>Effective</i></td> <td></td> </tr> </table>	<i>Effective</i>		<table border="1"> <tr> <td><i>Neutral</i></td> <td>X</td> </tr> </table>	<i>Neutral</i>	X	<table border="1"> <tr> <td><i>Ineffective</i></td> <td></td> </tr> </table>	<i>Ineffective</i>		<table border="1"> <tr> <td><i>Unknown</i></td> <td></td> </tr> </table>	<i>Unknown</i>			
<i>Effectiveness of measure</i>																								
<i>Effective</i>																								
<i>Neutral</i>	X																							
<i>Ineffective</i>																								
<i>Unknown</i>																								
<p><i>Rationale:</i> Macrophyte mapping by professional scuba divers is commonly used in EU countries in accordance to the WFD. The presence of a target species within the studied transects is determined with high accuracy, but infestations outside the transects in the study area will not be detected.</p>																								
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>Macrophyte mapping of evergreen species like <i>L. major</i> can be carried out throughout the year. In lakes with dominant native vegetation, when e.g. small <i>L. major</i> stands are embedded in dense beds of native macrophytes during the summer period, macrophyte mapping during winter time can help to identify small patches of this evergreen invasive alien aquatic plant species (author's observation).</p>																							
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Macrophyte mapping by scuba divers requires specialised diving equipment and some experience in submerged macrophyte mapping, independently if the investigations are carried out by professional or citizen scientist divers. However, by developing identification keys for the public and apps for mobiles (see citizen science table above), the cost of monitoring can be reduced and larger areas can be surveyed.</p> <p>Even though the costs of mapping invasive aquatic plants are hard to quantify, it should be borne in mind that the surveillance of submerged plants in aquatic habitats is much more time-consuming and costly than mapping plants in terrestrial habitats.</p>																							
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	<i>Environmental effects</i>							<table border="1"> <tr> <td><i>Positive</i></td> <td>X</td> </tr> </table>	<i>Positive</i>	X	<table border="1"> <tr> <td><i>Neutral or mixed</i></td> <td></td> </tr> </table>	<i>Neutral or mixed</i>		<table border="1"> <tr> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Negative</i>								
<i>Environmental effects</i>																								
<i>Positive</i>	X																							
<i>Neutral or mixed</i>																								
<i>Negative</i>																								
		<table border="1"> <tr> <td><i>Social effects</i></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	<i>Social effects</i>						<table border="1"> <tr> <td><i>Positive</i></td> <td></td> </tr> </table>	<i>Positive</i>		<table border="1"> <tr> <td><i>Neutral or mixed</i></td> <td>X</td> </tr> </table>	<i>Neutral or mixed</i>	X	<table border="1"> <tr> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Negative</i>								
<i>Social effects</i>																								
<i>Positive</i>																								
<i>Neutral or mixed</i>	X																							
<i>Negative</i>																								
		<table border="1"> <tr> <td><i>Economic effects</i></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>	<i>Economic effects</i>						<table border="1"> <tr> <td><i>Positive</i></td> <td></td> </tr> </table>	<i>Positive</i>		<table border="1"> <tr> <td><i>Neutral or mixed</i></td> <td>X</td> </tr> </table>	<i>Neutral or mixed</i>	X	<table border="1"> <tr> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Negative</i>								
<i>Economic effects</i>																								
<i>Positive</i>																								
<i>Neutral or mixed</i>	X																							
<i>Negative</i>																								
<p><i>Rationale:</i> During macrophyte mapping by diving or using a sub-camera other IAAPs can be identified, which reduces the total cost of monitoring per species.</p>																								

<p>rationale, with supporting evidence and examples if possible.</p>									
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p><i>Acceptable</i></p>	<p>X</p>	<p><i>Neutral or mixed</i></p>		<p><i>Unacceptable</i></p>			
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>		<p>There is no information available.</p>							
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>		<p><i>Unresolved</i></p>	<p><i>Established but incomplete</i></p>	<p>X</p>	<p><i>Well established</i></p>			
<p><i>Rationale:</i> Repeated macrophyte mappings are carried out to map and evaluate the macrophyte community in accordance to the EU WFD in rivers and lakes >50 ha, and during these investigations infestations of invasive submerged plants are often found (Van de Weyer <i>et al.</i>, 2007). However, the use of this method as an early detection mechanism has not been studied.</p>									

Rapid eradication for new introductions - Measures to achieve eradication at an early stage of invasion, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the eradication measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Physical control - Suction dredging</p> <p>For the rapid eradication of early infestations of invasive submerged aquatic plant species, physical control measures are the mostly used eradication measures. Suction dredging, hand weeding and shading of small areas by benthic barriers, e.g. jute matting, are documented as successful eradication measures for early infestations of submerged plants like <i>L. major</i>. These measures can be used alone or in combination, or as follow-ups of other eradication and management measures. Each of these measures is addressed in a separate rapid eradication table.</p> <p>During suction dredging, the target plants, surrounding water and sediment are sucked up, and the plants are collected in a fine mesh bag on a floating barge (de Winton <i>et al.</i>, 2013). The level of control depends on the depth of sediment removed, as the whole root system must be removed for effective eradication. Due to the removal of the upper sediment, the rehabilitated sediment can be less suitable for fast and large reinfestations, which can provide effective control of <i>L. major</i> for up to three years (Wells <i>et al.</i>, 2000 in de Winton <i>et al.</i>, 2003). In combination with follow-up hand removal, submerged aquatic plants can be eradicated (de Winton <i>et al.</i>, 2013).</p>							
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Suction dredging is suitable for small areas (<0.1 ha) infested with the invasive submerged plant species (de Winton <i>et al.</i>, 2013), and is used in public amenity areas to reduce the risk of overland dispersal of aquatic weeds attached to water-sport equipment.</p>							
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>X</p>	<p><i>Neutral</i></p>	<p><i>Ineffective</i></p>	<p><i>Unknown</i></p>		
<p><i>Rationale:</i> Suction dredging results in effective control of <i>L. major</i> for up to three years (Wells <i>et al.</i>, 2000 in de Winton <i>et al.</i>, 2013) but, when used in combination with follow-up hand weeding, suction dredging has eradicated weeds from some sites in New Zealand lakes (de Winton <i>et al.</i>, 2013). In New Zealand, suction dredging is used to eradicate outlier colonies and new incursions of <i>L. major</i> (Wells & Clayton, 2005 in de Winton <i>et al.</i>, 2013), and it can have a small and short-term impact on extensive <i>L. major</i> infestations (Howard-Williams & Reid, 1989 in Clayton, 1996). Moreover, suction dredging is used to control aquatic weeds in public amenity areas, in order to reduce the risk of</p>								

	<p>overland dispersal of aquatic plant fragments attached to watersport equipment, such as boats and trailers (de Winton <i>et al.</i>, 2013). In Texas, suction dredging successfully eradicated a submerged plant from a river section (Alexander <i>et al.</i>, 2008).</p>						
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>Suction dredging can be best carried out in good water clarity, if the sediment has a fine or muddy structure (de Winton <i>et al.</i>, 2013). Suction dredging is time consuming, as only small areas can be controlled within a certain amount of time. Nevertheless, the overall costs are still low to moderate, as no maintenance is required, but a monitoring should be carried out on an annual basis (de Winton <i>et al.</i>, 2013). In some cases, hand-weeding must follow suction dredging to eradicate the remaining individual plants.</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Suction dredging requires a large suction pump and skilled diving operators. While for small ponds and infestations close to the shore vehicular access is required, a barge must be used in larger systems (de Winton <i>et al.</i>, 2013). In dense weed beds, suction dredging requires 20 days per ha (de Winton <i>et al.</i>, 2013). In New Zealand, the costs of suction dredging were between 5,000 to 7,500 EUR per ha in Lake Wanaka and between 10,000 to 13,000 EUR in Rotorua Lakes (Wells <i>et al.</i>, 2000 in de Winton <i>et al.</i>, 2013).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Environmental effects</i></p>	<p><i>Positive</i></p>		<p><i>Neutral or mixed</i></p>		<p><i>Negative</i></p>	<p>X</p>
	<p><i>Social effects</i></p>	<p><i>Positive</i></p>		<p><i>Neutral or mixed</i></p>		<p><i>Negative</i></p>	<p>X</p>
	<p><i>Economic effects</i></p>	<p><i>Positive</i></p>		<p><i>Neutral or mixed</i></p>	<p>X</p>	<p><i>Negative</i></p>	
	<p><i>Rationale:</i> Based on the experience of the diver, suction dredging can be highly species specific, consequently only having a minor impact on non-targeted plant species, although affecting the sediment-based biota (e.g. macroinvertebrates). While this measure is underway, invertebrates within the IAAPs and the upper sediment will be displaced, but direct effects on fish are unlikely as they can use avoidance behaviours to escape. The overall impact on the environment is low, and no impacts on public health are reported. Suction dredging can, however, temporarily impact the recreational use of water bodies, disturbing activities such as diving, as it increases turbidity within the water column. However, the increase in turbidity is lower than for other management measures (e.g. Hydro Venturi) (de Winton <i>et al.</i>, 2013).</p>						

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 151 1883 220"> <tr> <td>Acceptability to stakeholders</td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td><i>X</i></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> The acceptance of rapid eradication measures depends on the side effects of the measure and the knowledge about the potential ecological and environmental impact that large infestations of the target species might have. Suction dredging has an impact, although minor, on recreational activities performed in water bodies, so it might not be well accepted by some stakeholders.</p>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Unacceptable</i>		
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	<i>X</i>	<i>Unacceptable</i>				
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Rapid eradication measures have a high cost-effectiveness, as rapid eradication of early infestations is in general cheaper than later species management and control (Hussner <i>et al.</i>, 2017). The cost of inaction is hard to quantify but will be high, as <i>L. major</i> is suitable to grow in most water bodies within the EU.</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="640 858 1924 927"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td></td> <td><i>Well established</i></td> <td><i>X</i></td> </tr> </table> <p><i>Rationale:</i> Suction dredging is a well established tool for the eradication of early infestations of submerged aquatic weeds.</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	<i>X</i>
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	<i>X</i>		

Rapid eradication for new introductions - Measures to achieve eradication at an early stage of invasion, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the eradication measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Physical control - Hand weeding</p> <p>The removal of aquatic plants by hand is one of the most selective removal methods and can be used for small, localised infestations and for infestations within native plant communities that should not be negatively affected (de Winton <i>et al.</i>, 2013; Hussner <i>et al.</i>, 2016, 2017). During the removal, all plant parts, including their roots, should be removed to avoid any regrowth of the plants. Hand removal is suitable for small and limited incursions of submerged aquatic plant species (and is additionally used as a follow-up measure of other management measures at the larger scale). The removal by hand in shallow water is possible by wading, while in deeper water (>1.5 m) snorkelling or scuba diving is required (Bellaud, 2009).</p>								
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Hand weeding can be successfully used for small scale eradications, and works best if the target species can be easily identified within native macrophyte communities (requiring sufficient water clarity) and is only present at low densities of <125 shoots per 0.1 ha (Bellaud, 2009), in small (<1 m²) monospecific plant patches. In Lough Corrib, Ireland, a number of small and isolated <i>L. major</i> stands were eradicated by hand weeding carried out by divers (Caffrey <i>et al.</i>, 2009).</p> <p>In combination with manual raking and hand application of Hydro-Venturi (see management table below), approximately 45 m³ of <i>Myriophyllum aquaticum</i> were removed by hand from a small pond system (approximately 1 ha; Hussner, unpublished).</p>								
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Effectiveness of measures</p>	<p>Effective</p>	<p>X</p>	<p>Neutral</p>		<p>Ineffective</p>		<p>Unknown</p>	
<p>Effort required</p>	<p>The majority of plants (> 90% of the biomass of the target species; Hussner <i>et al.</i>, 2016) can be removed during the first hand weeding operation. Plant regrowth usually occurs and requires follow-up treatments, until all plant material of the target species has been successfully removed (de Winton <i>et al.</i>, 2013; Hussner <i>et al.</i>, 2016). De</p>								

e.g. period of time over which measure needs to be applied to have results	Winton <i>et al.</i> (2013) recommended monitoring for 3-5 years after removal of the last fragments before eradication of the species can be confirmed.						
Resources required ¹ e.g. cost, staff, equipment etc.	Hand weeding requires waders and snorkel or diving equipment. As for all working activities in aquatic systems, human safety requires, among others, personal floatation devices and skills in working in aquatic systems (e.g. boat handling and diving; de Winton <i>et al.</i> , 2013). De Winton <i>et al.</i> (2013) estimated the costs of two hand weeding treatments to achieve weed eradication at about 12,000 EUR per ha. The costs of hand weeding in combination with raking and Hydro-Venturi, for the eradication of 45 m ³ of <i>Myriophyllum aquaticum</i> , were ca. 18,000 EUR (Hussner, unpublished).						
Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc. For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	Environmental effects	Positive		Neutral or mixed	X	Negative	
	Social effects	Positive		Neutral or mixed	X	Negative	
	Economic effects	Positive		Neutral or mixed	X	Negative	
	<i>Rationale:</i> Hand weeding is one of the most species specific control measures, with minimal impact on native plants (Hussner <i>et al.</i> , 2016), as long as the hand weeding is carried out by skilled workers. During hand weeding and uprooting of plants, an increase in turbidity of the water might occur, but this has no sustained effect on the environment.						
Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc. Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	Acceptability to stakeholders	Acceptable	X	Neutral or mixed		Unacceptable	
	<i>Rationale:</i> The acceptance of rapid eradication measures depends on the side effects of the measure and the knowledge about the potential ecological and environmental impact that large infestations of the target species might have. As hand weeding has no effect on economic activities, and only a very minor and temporary negative impact on the ecosystem, a high acceptance of this measure by stakeholders and the public is highly likely.						
Additional cost information ¹ When not already included above, or in the species Risk Assessment.	Rapid eradication measures have a high cost-effectiveness, as rapid eradication of early infestations is in general cheaper than later species management and control (Hussner <i>et al.</i> , 2017). The cost of inaction is hard to quantify but will be high, as <i>L. major</i> is suitable to grow in most water bodies within the EU.						

<ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>									
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	<i>X</i>	<p><i>Rationale:</i> Hand weeding is well established as a tool for the eradication of early infestations of submerged aquatic weeds.</p>

<p>Rapid eradication for new introductions - Measures to achieve eradication <u>at an early stage of invasion</u>, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the eradication measures identified.</p>	
<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Physical control - Benthic barriers</p> <p>Benthic barriers shade the sediment, hindering the plants to root in the sediment (de Winton <i>et al.</i>, 2013; Gettys <i>et al.</i>, 2014). Benthic barriers are used for the control of submerged aquatic plants (Boylen <i>et al.</i>, 1996; Caffrey <i>et al.</i>, 2010; Laitala <i>et al.</i>, 2012; Hoffmann <i>et al.</i>, 2013), but their use is limited to stagnant waters. While in the past plastic sheets were used, in recent years biodegradable jute, hessian or coconut mattings have been tested, with jute and coconut mats showing the best results (Caffrey <i>et al.</i>, 2010; Hofstra & Clayton, 2012). In contrast to plastic or polyethylene, loose woven biodegradable mats like jute mats allow the exchange of water and gas, allow some native plants to grow through the mats (Caffrey <i>et al.</i>, 2010; Hoffmann <i>et al.</i>, 2013) and allow native plant communities to regrow after the natural degradation of the material (Caffrey <i>et al.</i>, 2010; de Winton <i>et al.</i>, 2013).</p>

	<p>Benthic barriers made of biodegradable material can be used in small ponds and lakes, when only small infestations (<0.4 ha; de Winton <i>et al.</i>, 2013) of the target species are present, and if only temporary installation of benthic barriers is required (de Winton <i>et al.</i>, 2013), while for long-term control treatments, long-lasting polypropylene mats are the best option (de Winton <i>et al.</i>, 2013).</p> <p>It is important to note that high suspended sediment loading in the water column causes an accumulation of sediment on top of the benthic barriers, which provides a suitable substrate for new infestations of both native and invasive aquatic plants on the barriers (de Winton <i>et al.</i>, 2013; Hoffmann <i>et al.</i>, 2013).</p>						
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Benthic barriers have been used for infestations up to 5,000 m² (Caffrey <i>et al.</i>, 2010; de Winton <i>et al.</i>, 2013). In Lough Corrib, Ireland, a total area of 1,725 m² infested with <i>L. major</i> was covered with jute mats in 2008 (Caffrey <i>et al.</i>, 2009).</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 751 1906 818"> <tr> <td data-bbox="645 751 927 818"><i>Effectiveness of measures</i></td> <td data-bbox="927 751 1135 818"><i>Effective</i></td> <td data-bbox="1135 751 1191 818">X</td> <td data-bbox="1191 751 1377 818"><i>Neutral</i></td> <td data-bbox="1377 751 1624 818"><i>Ineffective</i></td> <td data-bbox="1624 751 1850 818"><i>Unknown</i></td> </tr> </table> <p><i>Rationale:</i> Benthic barriers are effective in stagnant waters, and have been shown to cause decomposition of <i>L. major</i> in Lough Corrib, Ireland (Caffrey <i>et al.</i>, 2009, 2010).</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>	<i>Ineffective</i>	<i>Unknown</i>
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>	<i>Ineffective</i>	<i>Unknown</i>		
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>In Lough Corrib, four months of shading by jute mattings resulted in the decomposition of <i>L. major</i> (Caffrey <i>et al.</i>, 2010).</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Most of the effort is required during the installation of the benthic barriers. Rolls of the benthic barrier (usually 5 m in width; Caffrey <i>et al.</i>, 2010) are used, and the material is placed onto the water surface from a boat-mounted dispenser. While plastic is difficult to sink and commonly floats up due to the gas production resulting from decaying plant material beneath it, biodegradable mattings made out of jute rapidly saturate with water and sink within minutes (Caffrey <i>et al.</i>, 2010), not floating up due to the loose-woven structure which allows gas exchange (Caffrey <i>et al.</i>, 2010). The installation of benthic barriers requires scuba divers to put the barriers in place and to anchor the</p>						

	<p>barriers within the sediment by using weights (rock or sand bags, or a layer of gravel or sand) or pins (Caffrey <i>et al.</i>, 2009, 2010; de Winton <i>et al.</i>, 2013; Gettys <i>et al.</i>, 2014). A reduction of the biomass may be necessary prior to the installation of the barriers, if submerged plants have already built up large stands which do not allow the barriers to be put in place. While plastic or polyethylene barriers must be removed from the habitat after the control measures, biodegradable mats can stay in place. The costs for benthic barriers are ca. 20,000 EUR per ha (one treatment, de Winton <i>et al.</i>, 2013).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>Negative</p>	<p>X</p>	
	<p>Social effects</p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>X</p>	<p>Negative</p>	
	<p>Economic effects</p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>X</p>	<p>Negative</p>	
	<p><i>Rationale:</i> Benthic barriers, particularly those made of plastic or polyethylene, affect all organisms beneath the barrier, including algae and submerged plants, as well as macroinvertebrates and benthic fish. However, if loose-woven biodegradable material like jute is used, native plants (e.g. charophytes) can grow through the barriers, resulting in a rapid recovery of native plant communities (Caffrey <i>et al.</i>, 2010).</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>		<p>Acceptable</p>	<p>X</p>	<p>Neutral or mixed</p>	<p>Unacceptable</p>	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness</p>	<p>Rapid eradication measures have a high cost-effectiveness, as rapid eradication of early infestations is in general cheaper than later species management and control (Hussner <i>et al.</i>, 2017). The cost of inaction is hard to quantify but will be high, as <i>L. major</i> is suitable to grow in most water bodies within the EU.</p>						

<p>- the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	<i>X</i>	<i>Well established</i>	
	<p><i>Rationale:</i></p> <p>Benthic barriers are established tools for the eradication of early infestations of submerged aquatic weeds.</p>							

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Mowing and cutting</p> <p>Mowing and cutting is the most common method to manage large infestations of submerged aquatic weeds in Europe (Hussner <i>et al.</i>, 2017). Cutting and mowing boats are used either with or without subsequent harvest of the cut biomass (Gettys <i>et al.</i>, 2014; Hussner <i>et al.</i>, 2017; Kuiper <i>et al.</i>, 2017).</p> <p>While the cutting depth of mowing boats is often limited to a maximum of 2 m (although some cutter boats in the USA can cut up to 5 m in depth; Podraza <i>et al.</i>, 2008; Caffrey <i>et al.</i>, 2009; de Winton & Clayton, 2016; Hussner <i>et al.</i>, 2017), V-blades can be pulled along the lake bed and rip plants from the substrate, resulting in an uprooting of plants and causing greater damage than cutting in the upper parts of the plants (Caffrey <i>et al.</i>, 2011). The efficiency of the cutting can be influenced by the morphology of <i>L. major</i> during the measure application (Caffrey <i>et al.</i>, 2009). In Lough Corrib, Ireland, two different morphologies of <i>L. major</i> were found, with a “collapsed” form between May and October and an “erect” form between October and March, being the latter form more easy to cut and harvest (Caffrey <i>et al.</i>, 2009).</p>

<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Mowing and cutting is mostly used as a control measure for infestations of submerged weeds in large scales, like it has been done for <i>Elodea nuttallii</i> in large lakes and reservoirs in Germany (Podraza <i>et al.</i>, 2008; Zehnsdorf <i>et al.</i>, 2015). In Ireland, about 29.2 ha of <i>L. major</i> infestations were cut and about 4,700 tons of cut plant material was subsequently harvested (Caffrey <i>et al.</i>, 2009).</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<input type="checkbox"/>	<p><i>Neutral</i></p>	<input checked="" type="checkbox"/>	<p><i>Ineffective</i></p>	<input type="checkbox"/>	<p><i>Unknown</i></p>	<input type="checkbox"/>	
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>All weed cutting and mowing measures must be repeated up to several times within a year to produce a significant reduction in biomass of the target species. Cutting by using V-blades is reported to be more effective than mowing, allowing longer intervals between treatments.</p>									
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>For mowing and cutting, boats and harvesters are required, as well as skilled operators. De Winton <i>et al.</i> (2013) noted costs of 2,000-4,000 NZD (ca. 1,175–2,350 EUR) per ha for mowing and harvesting the plant material, with</p>									

	costs varying with the distance to the dump site to deposit the harvested material. The costs for mowing and harvesting submerged <i>Elodea nuttallii</i> were about 2,500 EUR per day for 0.5 ha (Podraza, 2017).																										
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> </table>	Environmental effects	Positive		Neutral or mixed		Negative	X	Social effects	Positive		Neutral or mixed	X	Negative		Economic effects	Positive		Neutral or mixed	X	Negative		<p><i>Rationale:</i> Mechanical harvesting is commonly considered as having no impact on the environment, but fish and macroinvertebrates might be entrapped and killed within the harvested biomass (de Winton & Clayton, 2016). Furthermore, mowing and cutting are not species specific, and thus native plants localised within the treated area will be affected as well (Hussner <i>et al.</i>, 2017). Moreover, it must be considered that mowing and cutting produce large amounts of plant fragments, which can lead to an even faster spread of the target species (Anderson, 1998). The turbidity of the water column might also increase due to the uprooting of plants and the V-blade pulled along the sediment surface. Harvesting large amounts of biomass reduces the nutrient pool of aquatic systems (Kuiper <i>et al.</i>, 2017), but this is the case for all management methods which remove biomass.</p>				
Environmental effects	Positive		Neutral or mixed		Negative	X																					
Social effects	Positive		Neutral or mixed	X	Negative																						
Economic effects	Positive		Neutral or mixed	X	Negative																						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td>Acceptable</td> <td>X</td> <td>Neutral or mixed</td> <td></td> <td>Unacceptable</td> <td></td> </tr> </table>	Acceptability to stakeholders	Acceptable	X	Neutral or mixed		Unacceptable		<p><i>Rationale:</i> The public perception of mowing and cutting is positive, as this is usually carried out when large weed beds prohibit the recreational use of water bodies.</p>																		
Acceptability to stakeholders	Acceptable	X	Neutral or mixed		Unacceptable																						
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p>	<p>Mowing and cutting have not been documented as providing long-term effects in eradicating a submerged aquatic weed. While mowing in water depth up to 2 m does not lead to a substantial decrease in the biomass of the target species, V-blades which are pulled behind a boat across the sediment surface can significantly decrease the abundance of the target species within the treated area (Caffrey <i>et al.</i>, 2011). Thus, the cost-effectiveness for mowing must be considered low, while the cost-effectiveness for cutting along the sediment using V-blades shows a higher efficiency.</p>																										

Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).	However, the cost of inaction is high, as canopy forming submerged weeds like <i>L. major</i> can hinder the recreational use of water bodies, which might result in decreasing values of lakefront properties, as it has been documented for lakes infested with <i>Myriophyllum heterophyllum</i> (Halstead <i>et al.</i> , 2003).							
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
<p><i>Rationale:</i></p> <p>There are several studies documenting the effects of mowing and cutting with V-blades on <i>L. major</i> infestations, but meta-analyses are lacking.</p>								

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Hydro-Venturi</p> <p>The hydro-venturi (water jet) system pumps water into the upper sediment and washes the rooted plants out, which then float up and can be harvested. As whole plants are uprooted, there is less regrowth and less fragments are produced in comparison to other mechanical management measures, like mowing and cutting (van Valkenburg, 2011; Dorenbosch & Bergsma, 2014; Hussner <i>et al.</i>, 2017). This system has not yet been applied to <i>L. major</i>, but is expected to be suitable, based on its use on other invasive alien aquatic plants.</p> <p>The hydro-venturi works best in soft sediments (van Valkenburg, 2011; Hussner, unpublished) and has so far only been used in up to 1.5 m of water depth (van Valkenburg, 2011; Podraza, 2017), although its use in deeper water is possible. It is not species specific, but skilled operators can limit the impact on other aquatic plants.</p>
<p>Scale of application</p>	<p>The hydro-venturi was used for the control of <i>Cabomba caroliniana</i> and <i>Myriophyllum heterophyllum</i> in a shallow lake in the Netherlands (van Valkenburg, 2011). In combination with hand weeding, the application of hydro-venturi</p>

<p>At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>was used to eradicate approximately 45 m³ of <i>Myriophyllum aquaticum</i> in two connected ponds (total area about 1 ha), which resulted in > 99% reduction of the biomass of the target species, and only approximately 15 L of plants were found during the first post-treatment, three months after the management measures (Hussner, unpublished).</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>X</p>	<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>		<p><i>Unknown</i></p>		
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p><i>Rationale:</i> While not yet used on <i>L. major</i>, it is expected that this system would be as effective to reduce (control) the species impacts, as it has been shown to be for other invasive alien aquatic plants. It was successfully used for the reduction of biomass and abundance of evergreen <i>Cabomba caroliniana</i> and <i>Myriophyllum heterophyllum</i> in shallow water systems in the Netherlands (van Valkenburg, 2011) and Germany (Hussner, unpublished). A hand application of the hydro-venturi system was successfully tested to reduce the biomass of submerged <i>Myriophyllum heterophyllum</i> in shallow waters with a maximum depth of approximately 50 cm: > 99% of biomass reduction was achieved after a single treatment, controlled one year after the treatment (Hussner, unpublished). In reservoirs of the River Ruhr, Germany, hydro-venturi control of <i>Elodea nuttallii</i> was not successful (Podraza, 2017), but no reasons for this failure of the control measures with hydro-venturi are noted.</p> <p>The efficiency of the control by using hydro-venturi depends on the sediment structure and water depth, as the water jet must be pumped into the upper sediment layer for successful uprooting of the plants.</p>									
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The operation of the hydro-venturi system requires 1-2 skilled operators, and the subsequent harvest of the uprooted plant material requires additional workers. The cost of the hydro-venturi system operated by boat is ca. 1.35-2.05 Euro/m² (depending on sediment structure and other variables), while the cost for management using hand application is ca. 5 Euro/m² (Hussner, unpublished).</p>									

<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> <td></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> <td></td> <td></td> </tr> </table> <p><i>Rationale:</i> The hydro-venturi washes the rooted plants out of the sediment, and thus an increase of the water turbidity during the control measures will occur. Any organisms within the sediment are dislocated by the water jet, and macroinvertebrates and fish might be entrapped within the harvested plant material (like for all management measures which harvest plant material out of the water). Hydro-venturi is not species specific, and thus effects on other plant species seem likely, although this negative impact can be reduced if the hydro-venturi is carried out by skilled operators. In comparison to mowing and cutting, hydro-venturi produces less fragments, reducing the risk for subsequent spread of the species being controlled.</p>								<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X			<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>				<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>			
	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																												
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																														
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																														
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> <td></td> <td></td> </tr> </table> <p><i>Rationale:</i> As for all other management methods, the control of invasive aquatic plants by hydro-venturi has a high level of acceptability by stakeholders and the public, particularly in areas where the nuisance growth of aquatic weeds like <i>L. major</i> prohibits the recreational use of the water. Despite the disturbance of aquatic fauna, such as macroinvertebrates, and other non-target plant species, no negative effects on native animals are reported.</p>								<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																					
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																														
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As for all management methods, the cost of inaction is usually high and will result in spreading of the target species, reducing the likelihood of future eradication and increasing management costs. The cost of inaction is also high, as canopy forming submerged weeds like <i>L. major</i> can hinder the recreational use of water bodies, which might result in decreasing values of lakefront properties, as it has been documented for lakes infested with <i>Myriophyllum heterophyllum</i> (Halstead <i>et al.</i>, 2003).</p>																																		
<p>Level of confidence on the information provided ²</p>	<table border="1"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td>X</td> <td><i>Established but incomplete</i></td> <td></td> <td><i>Well established</i></td> <td></td> <td></td> </tr> </table>								<i>Inconclusive</i>		<i>Unresolved</i>	X	<i>Established but incomplete</i>		<i>Well established</i>																				
<i>Inconclusive</i>		<i>Unresolved</i>	X	<i>Established but incomplete</i>		<i>Well established</i>																													

<p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Rationale:</i></p> <p>Hydro-venturi has been successfully used to control some aquatic weed species in shallow waters, but the control of <i>Elodea nuttallii</i> in the reservoirs of the River Ruhr was not successful. More research is needed to improve the success and applicability of this system. In addition, hydro-venturi has not been tested for the control of <i>L. major</i> so far (although it seems reasonable that it will be a valuable tool for the control of the species).</p>
--	--

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Biological control</p> <p>Biological control agents can be used to reduce the growth or reproductive capacity of a target species (Cuda <i>et al.</i>, 2008). There are various types of biocontrol, including generalist herbivores, the inundative and the classical and augmentative biocontrol (Hussner <i>et al.</i>, 2017).</p> <p>Generalist herbivores can be either native or non-native species which have a broad host range. The species are introduced with the aim to control the target plants. For the control of submerged aquatic weeds like <i>L. major</i>, grass carp (<i>Ctenopharyngodon idella</i>) is considered as a potential generalist control agent (Chapman & Coffey, 1971) and particularly a sterile triploid form is widely used (Venter & Schoonbee, 1991); however, grass carp is itself an invasive alien species in Europe. Grass carp have been stocked in a number of European lakes to control other submerged weeds like <i>Elodea nuttallii</i>. This often resulted in a decrease of all submerged plant species in the water body, as grass carp are not species specific (Dibble & Kovalenko, 2009; Hussner <i>et al.</i>, 2017).</p> <p>It should be borne in mind that the release of macro-organisms as biological control agents is currently not regulated at EU level. Nevertheless, national/regional laws are to be respected. Before any release of an alien species as a biological control agent, an appropriate risk assessment should be made.</p> <p>For classical biological control, the potential biological control agent is collected within the native range and introduced into the invasive range of the target species (van Driesche <i>et al.</i>, 2010). Host specificity tests are required prior to the release of the control agent to ensure that host shift will not occur. While for floating and emergent</p>

	<p>aquatic plants several successful examples of classical biological control are documented, there are no examples of successful classical biological control of submerged weeds (Hussner <i>et al.</i>, 2017). Several phytophagous insects (including e.g. <i>Bagous</i> sp., <i>Nymphulinae</i> sp., <i>Polypedilum</i> n.sp.) have been documented feeding on <i>L. major</i>, some of which have a high potential as biocontrol agents of the species (Baars <i>et al.</i>, 2010; Earle <i>et al.</i>, 2013). The most studied species for classical biological control of <i>L. major</i> is <i>Hydrellia lagarosiphon</i>, a leaf mining fly (Mangan & Baars, 2013; Martin <i>et al.</i>, 2013; Mangan <i>et al.</i>, 2019), but even for this species studies about the success to control <i>L. major</i> under field conditions are lacking.</p> <p>The inundative control of aquatic weeds using mycoherbicides is a new technique which was tested successfully in mesocosm experiments, but under field conditions no success in the control of submerged aquatic weeds is reported so far (Shearer, 1994; 1996; Hofstra <i>et al.</i>, 2004; Hussner <i>et al.</i>, 2017). There is no information on the use of mycoherbicides to control <i>L. major</i>.</p>									
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>No biological control agents have been released so far for the control of <i>L. major</i>.</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 858 1888 922"> <tr> <td data-bbox="645 858 925 922">Effectiveness of measures</td> <td data-bbox="925 858 1115 922">Effective</td> <td data-bbox="1115 858 1171 922"></td> <td data-bbox="1171 858 1377 922">Neutral</td> <td data-bbox="1377 858 1433 922"></td> <td data-bbox="1433 858 1624 922">Ineffective</td> <td data-bbox="1624 858 1680 922"></td> <td data-bbox="1680 858 1832 922">Unknown</td> <td data-bbox="1832 858 1888 922">X</td> </tr> </table> <p><i>Rationale:</i> There is no information about the use and success of generalist, classical, or inundative biological control agents to control <i>L. major</i> in the field. However, grass carp has proven to be a successful control agent for other submerged aquatic weeds (de Winton <i>et al.</i>, 2013), showing their potential effectiveness in controlling <i>L. major</i>.</p>	Effectiveness of measures	Effective		Neutral		Ineffective		Unknown	X
Effectiveness of measures	Effective		Neutral		Ineffective		Unknown	X		
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>For control using generalist herbivores, grass carp must be stocked in appropriate quantities (Dibble & Kovalenko, 2009) for several years to prevent any regrowth from vegetative means in the sediment.</p> <p>For <i>L. major</i>, no classical or inundative biocontrol control agents have been tested under field conditions so far. However, in general, host specificity testing would take about 3 years prior to the potential release of a control agent (Newman & Duenas, 2017).</p>									

<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The cost of the use of grass carp depends on the stocking densities needed. High costs will be incurred for the successful removal of grass carp after the control of the target species, to allow for the restoration of native plant communities (Hussner <i>et al.</i>, 2017).</p> <p>The testing of inundative and classical biological control agents requires experienced scientists and also involves high costs.</p>																										
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> </table>	Environmental effects	Positive		Neutral or mixed		Negative	X	Social effects	Positive		Neutral or mixed		Negative	X	Economic effects	Positive		Neutral or mixed	X	Negative				X		
Environmental effects	Positive		Neutral or mixed		Negative	X																					
Social effects	Positive		Neutral or mixed		Negative	X																					
Economic effects	Positive		Neutral or mixed	X	Negative																						
<p><i>Rationale:</i> The side effects are indicated only for the use of grass carps, which are proven successful control agents for submerged aquatic weeds (de Winton <i>et al.</i>, 2013). However, they are themselves an invasive alien species in Europe and consume all kind of aquatic plants, so impacts on native plants are likely to be high. The disappearance of aquatic vegetation may affect waterbirds, which use aquatic plants and associated macroinvertebrates as food. The disappearance of submerged aquatic plants can also cause a shift to a phytoplankton-dominated state in the ecosystem, with increased turbidity, which makes water bodies less attractive for recreational activities and tourism.</p> <p>The use of classical biological and inundative control agents has not been tested in the field so far for <i>L. major</i>, but Newman and Duenas (2017) noted that there should be no side effects on native plant species, if the classical biological control programme is well-managed.</p>																											
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td>Acceptable</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Unacceptable</td> <td></td> </tr> </table>	Acceptability to stakeholders	Acceptable		Neutral or mixed	X	Unacceptable				X																
Acceptability to stakeholders	Acceptable		Neutral or mixed	X	Unacceptable																						
<p><i>Rationale:</i> The potential negative effects caused by grass carps may reduce the acceptability of this measure by stakeholders, as turbid waters are less attractive for recreational activities and tourism in general.</p> <p>For potential inundative and classical biological control agents for <i>L. major</i>, there is a large knowledge gap about the impacts that these control agents might cause, and consequently their acceptability is hard to quantify.</p>																											
<p>Additional cost information¹</p>	<p>The cost-effectiveness is often considered high for grass carp use, if only the costs of stocking are considered (about 1,000 NZD (ca. 587 EUR) per ha; Clayton, 1996). However, the follow-up costs make stocking of grass carp an</p>																										

<p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>expensive management strategy. The costs can be reduced in water bodies where water-level drawdown can be used to remove the grass carp after treatment.</p> <p>No data about the costs of classical and inundative biological control are available, but both methods require comprehensive testing prior to their use, incurring high costs. The costs for the testing phase are ca. 300,000 EUR (Newman & Duenas, 2017), and its cost-effectiveness has been estimated to be from 2.5:1 to 15:1, and even up to 4000:1 (McConnachie <i>et al.</i>, 2003 and Culliney, 2005 in Newman & Duenas, 2017).</p> <p>As for all control strategies, the costs of inaction will be high, as <i>L. major</i> is able to grow in most water bodies within the EU member countries.</p>							
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p>X</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	
<p><i>Rationale:</i></p> <p>Generalist herbivores such as grass carps are widely used to control submerged weeds, in general, and have been suggested for the control of <i>L. major</i> (Chapman & Coffey, 1971), but information on their use and success is lacking.</p> <p>Potential classical inundative and biological control agents (like <i>Hydrellia lagarosiphon</i>) have been identified for <i>L. major</i>, but there are no data from field tests available.</p>								

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Herbicides</p> <p>Herbicides, in general terms, are used to control aquatic plants in ponds and lakes, channels and irrigation systems (de Winton <i>et al.</i>, 2013; Gettys <i>et al.</i>, 2014; Hussner <i>et al.</i>, 2017). Herbicide treatment may significantly reduce the biomass of submerged weeds like <i>L. major</i> and can result in the eradication of a target species (de Winton <i>et al.</i>, 2013; Champion & Wells, 2014; Hussner <i>et al.</i>, 2017). While herbicides are usually not species specific, the chosen</p>

	<p>concentration, exposure time, species specific uptake rates of a given herbicide and the application method used can cause a level of selectivity (Getsinger <i>et al.</i>, 1997, 2008, 2014; Carvalho <i>et al.</i>, 2007; Netherland, 2014).</p> <p>EU/national/local legislation on the use of plant protection products and biocides needs to be respected when applying this measure. According to Regulation (EC) No 1107/2009 concerning plant protection products, none of the active substances mentioned below are approved for use in the EU, although national authorisations might be possible. It is important to also comply with Regulation (EU) No 528/2012 on the use of biocidal products.</p>									
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Herbicides are commonly used for the large scale control of submerged aquatic weeds (Clayton, 1996). In Lake Rotorua, New Zealand, herbicide treatment was used to treat >100 ha of invasive aquatic weeds (Clayton, 1996). In Lough Corrib (Ireland), only parts of the lakes and a harbour were treated with the herbicide dichlobenil (Caffrey <i>et al.</i>, 2009).</p>									
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 679 1888 743"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> <td><i>Unknown</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> For <i>L. major</i>, some herbicides have been tested, with varying levels of control achieved (Clayton, 1996; Hofstra & Clayton, 2001; Caffrey <i>et al.</i>, 2009; Hussner <i>et al.</i>, 2017). Endothal and diquat, which are registered for use in New Zealand, have been tested to control <i>L. major</i> in the country (de Winton & Clayton, 2016) and successful treatments have caused a substantial reduction in the species biomass (Clayton, 1996; de Winton & Clayton, 2016). In Lough Corrib, Ireland, the application of dichlobenil granules did not result in a remarkable reduction of <i>L. major</i> coverage (Caffrey & Avedo, 2008), which was probably a result of the granules being trapped in the vegetation. Consequently, they did not reach the lake bed, which is the site of activity for dichlobenil (Caffrey, 1993a,b in Caffrey & Acevedo, 2008). However, in suitable habitats, a 100% eradication of <i>L. major</i> is normally achieved (Caffrey <i>et al.</i>, 2011).</p>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>		<i>Unknown</i>	
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>		<i>Unknown</i>			
<p>Effort required e.g. period of time over which measure needs to be applied to have results</p>	<p>The frequency and amount of herbicide needed for successful control depends on the herbicide and its formula used (Hofstra & Clayton, 2001; Hussner <i>et al.</i>, 2017). In Lough Corrib, a single treatment with dichlobenil resulted in total weed kill in suitable habitats (Caffrey <i>et al.</i>, 2011).</p>									
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Besides the herbicide, a boat and experienced workers are needed for the surface application of herbicides like dichlobenil. In water bodies with a high biomass of the target species, aerial application is possible (de Winton <i>et al.</i>, 2013).</p>									

	The full cost (herbicide, operator and equipment) of herbicide treatment with gel-formulated diquat in Lake Rotorua, New Zealand, was less than 350 EUR (500 NZD) per ha (Clayton, 1996).						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	Environmental effects	Positive		Neutral or mixed		Negative	X
	Social effects	Positive		Neutral or mixed		Negative	X
	Economic effects	Positive		Neutral or mixed	X	Negative	
	<p><i>Rationale:</i> In Lake Rotorua, diquat application enhanced the maintenance of native charophytes, which are resistant to this herbicide (Clayton, 1996). However, the effects of herbicides on other submerged aquatic plants depend on the sensitivity of the species to the herbicides, with varying effects at a species level. Besides the effects on other submerged plants, macroinvertebrates and fish can also be affected. Caffrey (1993a,b) described only minor negative effects of dichlobenil on water quality, non-target aquatic plants, macroinvertebrates and fish.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	Acceptability to stakeholders	Acceptable		Neutral or mixed	X	Unacceptable	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p><i>Rationale:</i> The chemical control of submerged aquatic weeds by using herbicides can be relatively cheap (Clayton, 1996), which might increase the acceptability of stakeholders, while the potential negative impact on other submerged plants, fauna and water quality could cause less acceptability. Until now, only dichlobenil was used for the control of <i>L. major</i> in Ireland (Caffrey <i>et al.</i>, 2009).</p> <p>While no specific information about the costs in the EU is available, the general low cost of aquatic weed control with herbicides (Clayton, 1996) makes its cost-effectiveness high. However, if the used herbicide negatively affects native plant communities, their restoration will have additional costs.</p>						

Level of confidence on the information provided ² Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure	<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input checked="" type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input type="checkbox"/>
	<i>Rationale:</i> The results from field trials with dichlobenil in Lough Corrib (Ireland) had varying results in the control of <i>L. major</i> (Caffrey <i>et al.</i> , 2009, 2011).							

Bibliography³

See guidance section

- Adriaens, T., Sutton-Croft, M., Owen, K., & Schneider, K., (2015). Trying to engage the crowd in recording invasive alien species in Europe: experiences from two smartphone applications in northwest Europe. *Management of Biological Invasions*, 6, 215–225.
- Alexander, M. L., Doyle, R. D., & Power, P. (2008). Suction dredge removal of an invasive macrophyte from a spring-fed river in Central Texas, USA. *Journal of Aquatic Plant Management*, 46, 184–185.
- Anderson, L. W. J. (1998). Dissipation and movement of Sonar and Komeen following typical applications for control of *Egeria densa* in the Sacramento–San Joaquin Delta and Production and viability of *E. densa* fragments following mechanical harvesting (1997/1998). Davis, CA U.S. Department of Agriculture–Agricultural Research Service, Environmental Impact Report for California Department of Boating and Waterways, 79pp.
- Anderson, L. W. J. (2005). California’s reaction to *Caulerpa taxifolia*: A Model for Invasive Species Rapid Response. *Biological Invasions*, 7, 1003-1016.
- Anderson, L. G., Dunn, A. M., Rosewarne, P. J., & Stebbing, P. D. (2015). Invaders in hot water: a simple decontamination method to prevent the accidental spread of aquatic invasive non-native species. *Biological Invasions*, 17, 2287–2297.
- Anderson, L. G., White, P. C., Stebbing, P. D., Stentiford, G. D., & Dunn, A. M. (2014). Biosecurity and vector behaviour: Evaluating the potential threat posed by anglers and canoeists as pathways for the spread of invasive non-native species and pathogens. *PLOS ONE*, 9(4), e92788.
- Baars, J. R., Coetzee, J. A., Martin, G., Hill, M. P., & Caffrey, J. M. (2010). Natural enemies from South Africa for biological control of *Lagarosiphon major* (Ridl.) Moss ex Wager (Hydrocharitaceae) in Europe. *Hydrobiologia*, 656, 149-158.
- Barnes, M. A., Jerde, C. L., Keller, D., Chadderton, W. L., Howeth, J. G., & Lodge, D. M. (2013). Viability of aquatic plant fragments following desiccation. *Invasive Plant Science and Management*, 6, 320-325.
- Bellaud, M. D. (2009). Cultural and physical control of aquatic weeds. Chapter 6. In: Gettys LA, Haller WT, Bellaud M (Eds). *Biology and control of aquatic plants, a best management practices handbook*. Aquatic Ecosystem Restoration Foundation, Georgia, USA.
- Boylen, C. W., Eichler, L. W., & Sutherland, J. W. (1996). Physical control of Eurasian watermilfoil in an oligotrophic lake. *Hydrobiologia*, 340, 213–218.
- Brunel, S. (2009). Pathway analysis: aquatic plants imported in 10 EPPO countries. *EPPO Bulletin*, 39, 201–213.
- Burchnall, W. (2013). Wetland Biosecurity Officer project report. The Broads Authority. http://www.broads-authority.gov.uk/__data/assets/pdf_file/0020/405344/Biosecurity_officer__6_month_update.pdf.
- Caffrey, J. M. (1993a). Aquatic weed management practices using dichlobenil: an Irish experience. *Polskie Archiwum Hydrobiologii*, 40, 255-26.
- Caffrey, J. M. (1993b). Plant management as an integral part of Ireland’s aquatic resources. *Hydroecologie Applique*, 5, 77-96.

- Caffrey, J. M., & Acevedo, S. (2008). *Lagarosiphon major* in Lough Corrib – Management options. In: C Moriary, R Rossell, & P Gargan (eds.). Fish stocks and their environment, 85-97. Westport, Institute of Fisheries Management.
- Caffrey, J. M., Evers, S., & Moran, H. (2009). Research and control programme for *Lagarosiphon major* in Lough Corrib 2008. Dublin Central Fisheries Board.
- Caffrey, J. M., Millane, M., Evers, S. L., & Moran, H. (2011). Management of *Lagarosiphon major* (Ridley) Moss in Lough Corrib – a review. *Biology and Environment: Proceedings of the Royal Irish Academy*, 111, 205-212.
- Caffrey, J. M., Millane, M., Evers, S., Moran, H., & Butler, M. (2010). A novel approach to aquatic weed control and habitat restoration using biodegradable jute matting. *Aquatic Invasions*, 5, 123-129.
- Carvalho, R. F. de, Bromilow, R. H., & Greenwood, R. (2007). Uptake of pesticides from water by curly waterweed *Lagarosiphon major* and lesser duckweed *Lemna minor*. *Pest Management Science*, 63, 789-797.
- Champion, P. D., & Clayton, J. S. (2001). Border Control for Potential Aquatic Weeds. Stage 2. Weed Risk Assessment. Science for Conservation 165. Department of Conservation, Wellington, New Zealand, 30pp.
- Champion, P. D., Clayton, J. S., & Hofstra, D. E. (2010). Nipping aquatic plant invasion in the bud: weed risk assessment and the trade. *Hydrobiologia*, 656, 167-172.
- Champion, P., & Wells, R. (2014). Proactive management of aquatic weeds to protect the nationally important Northland dune lakes. In: Nineteenth Australasian Weeds Conference, New Zealand. www.caws.org.au/awc/2014/awc201411391.
- Chapman, V. J., & Coffey, B. J. (1971). Experiments with grass carp in controlling exotic macrophytes in New Zealand. *Hydrobiologia*, 12, 313-323.
- Clayton, J. S. (1996). Aquatic weeds and their control in New Zealand lakes. *Journal of Lake and Reservoir Management*, 12, 477-486.
- Clayton, J. S., & Franklyn, G. (2005). Assessment of the December 2004 *Lagarosiphon major* control in Lake Wanaka. Internal Report, National Institute of Water and Atmospheric Research, New Zealand.
- Crane, K., Cuthbert, R. N., Dick, J. T. A., Kregting, L., MacIsaac, H. J., & Coughlan, N. E. (2019). Full steam ahead: direct steam exposure to inhibit spread of invasive aquatic macrophytes. *Biological Invasions*, 21, 1311-1321.
- Cuda, J. P., Charudattan, R., Grodowitz, M. J., Newman, R. M., Shearer, J. F., Tamayo, M. L., & Villegas, B. (2008). Recent advances in biological control of submersed aquatic weeds. *Journal of Aquatic Plant Management*, 46, 15-32.
- Culliney, T. W. (2005). Benefits of Classical Biological Control for managing invasive plants. *Critical Reviews in Plant Sciences*, 24, 131-150.
- Cuthbert, R. N., Coughlan, N. E., Crane, K., Caffrey, J. M., & MacIsaac, H. J. (2018). A dip or dab: assessing the efficacy of Virasure Aquatic disinfectant to reduce secondary spread of the invasive curly waterweed *Lagarosiphon major*. *Management of Biological Invasions*, 9, 259-265.

- Delaney, D. G., Sperling, C. D., Adams, C. S., & Leung, B. (2008). Marine invasive species: validation of citizen science and implications for national monitoring networks. *Biological Invasions*, 10, 117-128.
- de Winton, M., & Clayton, J. S. (2016). Ten year management plan for *Lagarosiphon major* at Lake Dunstan: 2016-2025. NIWA Client report No. HAM2016-040, 41pp.
- de Winton, M., Jones, H., Edwards, T., Özkundakci, D., Wells, R., McBride, C., Rowe, D., Hamilton, D., Clayton, J., Champion, P., & Hofstra, D. (2013). Review of best management practices for aquatic vegetation control in stormwater ponds, wetlands, and lakes. Prepared by NIWA and the University of Waikato for Auckland Council. Auckland Council technical report, TR2013/026.
- Dibble, E. D., & Kovalenko, K. (2009). Ecological impact of grass carp: a review of the available data. *Journal of Aquatic Plant Management*, 47, 1–15.
- Dorenbosch, M., & Bergsma, J. H. (2014). Bestrijding Van Waterwaaier in HardinxveldMet Hydro-venturi. Evaluatie Onderzoek En Beknopte Literatuur Vergelijking. Bureau Waardenburg, Culemborg.
- Earle, W., Mangan, R., O'Brien, M., & Baars, J. R. (2013). Biology of *Polypedilum* n.sp. (Diptera: Chironomidae), a promising candidate agent for the biological control of the aquatic weed *Lagarosiphon major* (Hydrocharitaceae) in Ireland. *Biocontrol Science and Technology*, 23, 1267-1283.
- Genovesi, P., Scalera, R., Brunel, S., Roy, D., & Solarz, W. (2010). Towards an early warning and information system for invasive alien species (IAS) threatening biodiversity in Europe. EEA Technical report No 5/2010, EEA, Copenhagen, 47pp.
- Getsinger, K. D., Turner, E. G., Madsen, J. D., & Netherland, M. (1997). Restoring native plant vegetation in a Eurasian watermilfoil dominated community using the herbicide triclopyr. *Regulated River*, 13, 357–375.
- Getsinger, K. D., Netherland, M. D., Grue, C. E., & Koschnick, T. J. (2008). Improvements in the use of aquatic herbicides and establishment of future research directions. *Journal of Aquatic Plant Management*, 46, 32–41.
- Getsinger, K. D., Dibble, E., Rodgers, J. H., & Spencer, D. (2014). Benefits of Controlling Nuisance Aquatic Plants and Algae in the United States. CAST Commentary, QTA2014-1, 12pp.
- Gettys, L., Haller, W., & Petty, D. (Eds.) (2014). Biology and Control of Aquatic Plants. A Best Management Practices Handbook: Third Edition. Aquatic Ecosystem Restoration Foundation, USA, 252pp.
- Halford, M., Heemers, L., van Wesemael, D., Mathys, C., Wallens, S., Branquart, E., Vanderhoeven, S., et al. (2014). The voluntary Code of conduct on invasive alien plants in Belgium: results and lessons learned from the AlterIAS LIFE+ project. *EPPO Bulletin*, 44, 1-11.
- Halstead, J. M., Michaud, J., Hallas-Burt, S., & Gibbs, J. P. (2003) Hedonic analysis of effects of a non-native invader (*Myriophyllum heterophyllum*) on a New Hampshire (USA) lakefront properties. *Environmental Management*, 32, 391-398.
- Heidbüchel, P., & Hussner, A. (2019). Fragment type and water depth determine the regeneration and colonization success of submerged aquatic plants. *Aquatic Sciences*, 81, 6.

- Heidbüchel, P., Jahns, P., & Hussner, A. (2019). Chlorophyll fluorometry sheds light on the role of desiccation resistance for vegetative overland dispersal of aquatic plants. *Freshwater Biology*, 64(8), 1401-1415.
- Hippolite, J., & Te Kurapa, T. W. M. (2018). Aquatic Pest Report 2018. Bay of Plenty Regional Council, Environmental Publication 2018/02, Whakatane, New Zealand.
- Hochachka, W. M., Fink, D., Hutchinson, R. A., Sheldon, D., Wong, W. K., & Kelling, S. (2012). Data intensive science applied to broad-scale citizen-science. *Trends in Ecology & Evolution*, 27, 130-137.
- Hoffmann, M., Gonzalez, A. B., Raeder, U., & Melzer, A. (2013). Experimental weed control of *Najas marina* ssp. *intermedia* and *Elodea nuttallii* in lakes using biodegradable jute matting. *Journal of Limnology*, 72, 485-493.
- Hofstra, D. E., & Clayton, J. S. (2012). Assessment of benthic barrier products for submerged aquatic weed control. *Journal of Aquatic Plant Management*, 50, 101-105.
- Hofstra, D., Shearer, J., Edwards, T., Clayton, J., & Gemmill, C. (2004). *Mycoleptodiscus terrestris*: A comparison between isolates from the United States and New Zealand. In: Aquatic Plant Management Society Conference, July 2004.
- Howard-Williams, C., & Reid, V. (1989). Aquatic weed survey of Lake Whakamarino following dredging. DSIR unpublished report to Electrocorp.
- Hussner, A. (2012). Alien aquatic plants in European countries. *Weed Research*, 52, 397-406.
- Hussner, A., Nehring, S., & Hilt, S. (2014). From first reports to successful control: a plea for improved management of alien aquatic plant species in Germany. *Hydrobiologia*, 737, 321-331.
- Hussner, A., Stiers, I., Verhofstad, M. J. J. M., Bakker, E. S., Grutters, B. M. C., Haury, J., van Valkenburg J. L. C. H., Brundu, G., Newman, J., Clayton, J. S., Anderson, L. W. J., & Hofstra, D. (2017). Management and control methods of invasive alien aquatic plants: a review. *Aquatic Botany*, 136, 113-137.
- Hussner, A., Windhaus, M., & Starfinger, U. (2016). From weed biology to successful control: an example of successful management of *Ludwigia grandiflora* in Germany. *Weed Research*, 56, 434-441.
- Johnson, L. E., Ricciardi, A., & Carlton, J. T. (2001). Overland dispersal of aquatic invasive species: a risk assessment of transient recreational boating. *Ecological Applications*, 11, 1789-1799.
- Johnstone, I. M., Coffey, B. T., & Howard-Williams, C. (1985). The role of recreational boat traffic in interlake dispersal of macrophytes: a New Zealand case study. *Journal of Environmental Management*, 20, 263-279.
- Kuiper, J. J., Verhofstad, M. J. J. M., Louwers, E. L. M., Bakker, E. S., Brederveld, R. J., van Gerven, L. P. A., Janssen, A. B. G., de Klein, J. J. M., & Mooij, W. M. (2017). Mowing submerged macrophytes in shallow lakes with alternative stable states: battling the good guys? *Environmental Management*, 59, 619-634.


- Laitala, K. L., Prather, T. S., Thill, D., Kennedy, B., & Caudill, C. (2012). Efficacy of benthic barriers as a control measure for Eurasian Watermilfoil (*Myriophyllum spicatum*). *Invasive Plant Science and Management*, 5: 170–177.
- Lockwood, J. L., Cassey, P., & Blackburn, T. (2005). The role of propagule pressure in explaining species invasions. *Trends in Ecology and Evolution*, 20, 223-228.
- Mangan, R., & Baars, J. R. (2013). Use of life table statistics and degree day values to predict the colonisation success of *Hydrellia lagarosiphon* Deeming (Diptera: Ephydriidae), a leaf mining fly of *Lagarosiphon major* (Ridley) Moss (Hydrocharitaceae), in Ireland and the rest of Europe. *Biological Control*, 64, 143-151.
- Mangan, R., Carolan, J. C., & Baars, J. R. (2019). Molecular characterization of *Hydrellia lagarosiphon*, a leaf mining biological control agent for *Lagarosiphon major*, reveals weak variance across large geographic areas in South Africa. *Biological Control*, 132, 8-15.
- Martin, G. D., Coetzee, J. A., & Baars, J. R. (2013). *Hydrellia lagarosiphon* Deeming (Diptera: Ephydriidae), a potential biological control agent for the submerged aquatic weed, *Lagarosiphon major* (Ridl.) Moss ex Wager (Hydrocharitaceae). *African Entomology*, 21, 151-160.
- Matsushashi, S., Doi, S., Fujiwara, A., Watanabe, S., & Minamoto, T. (2016). Evaluation of the Environmental DNA Method for Estimating Distribution and Biomass of Submerged Aquatic Plants. *PLoS One*, 11(6), e0156217.
- McConnachie, A. J., de Wit, M. P., Hill, M. P., & Byrne, M. J. (2003). Economic evaluation of the successful biological control of *Azolla filiculoides* in South Africa. *Biological Control*, 28, 25-32.
- Netherland, M. D. (2014). Chemical control of aquatic weeds. In: Gettys, L., Haller, W., Petty, D. (Eds.), *Biological Control of Aquatic Plants. A Best Management Practices Handbook*, third edition. Aquatic Ecosystem Restoration Foundation, USA, 71–88.
- Newman, J., & Duenas, M. A. (2017). Information on measures and related costs in relation to species included on the Union list: *Alternanthera philoxeroides*. Technical note prepared by IUCN for the European Commission.
- Podraza, P. (2017). Aquatische Makrophyten als Herausforderung für das Management von Flusstauseen am Beispiel der Ruhrstauseen. Oral presentation, AquaMak Conference, Leipzig, 30 March 2017.
- Podraza, P., Brinkmann, T., Evers, P., von Felde, D., Frost, U., Klopp, R., & Knotte, H. (2008). Untersuchungen zur Massenentwicklung von Wasserpflanzen in den Ruhrstauseen und Gegenmaßnahmen. Abschlussbericht des F and E-Vorhabens im Auftrag des Ministeriums für Umwelt und Naturschutz, Landwirtschaft und Verbraucherschutz des Landes NRW (MUNLV), 364 S.
- Redekop, P., Hofstra, D., & Hussner, A. (2016). *Elodea canadensis* shows a higher dispersal capacity via fragmentation than *Egeria densa* and *Lagarosiphon major*. *Aquatic Botany*, 130, 45-49.
- Roy, H. E., Pocock, M. J. O., Preston, C. D., Roy, D. B., Savage, J., Tweddle, J. C., & Robinson, L. D. (2012). Understanding citizen science & environmental monitoring. Final report on behalf of UK-EOF. NERC Centre for Ecology and Hydrology and Natural History Museum.

- Scriver, M., Marinich, A., Wilson, C., & Freeland, J. (2015). Development of species-specific environmental DNA (eDNA) markers for invasive aquatic plants. *Aquatic Botany*, 122, 27-31.
- Shearer, J. (1994). Field test of first commercial formulation of *Mycoleptodiscus terrestris* (Gerd.) Ostazeski as a biocontrol for Eurasian Watermilfoil. Miscellaneous. Paper A-94-4. US Army Waterways Experiment Station, Vicksbury, MS.
- Shearer, J. (1996). Field and laboratory studies of the fungus *Mycoleptodiscus terrestris* as a potential agent for management of the submerged aquatic macrophyte, *Hydrilla verticillate*. Technical report A-96-3. US Army Waterways Experiment Station, Vicksbury, MS.
- Van de Weyer, K., Nienhaus, I., Tigges, P., Hussner, A., & Hamann, U. (2007): Eine einfache und kosteneffiziente Methode zur flächenhaften Erfassung von submersen Pflanzenbeständen in Seen. *Wasser und Abfall*, 9, 20-22.
- Van de Wiel, C. C., van der Schoot, J., van Valkenburg, J. L. H. C., Duistermaat, H., & Smulders, M. J. (2009). DNA barcoding discriminates the noxious invasive plant species, floating pennywort (*Hydrocotyle ranunculoides* L.f.), from non-invasive relatives. *Molecular Ecology Resources*, 9, 1086-1091.
- Van Driesche, R. G., Carruthers, R. I., Center, T., Hoddle, M. S., Hough-Goldstein, J., Morin, L., Smith, L., et al. (2010). Classical biological control for the protection of natural ecosystems. *Biological Control*, 54, 2-33.
- van Valkenburg, J. L. H. C. (2011). *Cabomba caroliniana* and *Myriophyllum heterophyllum* a Nightmare Combination. Robson Meeting February 2011. www.robsonmeeting.org/valkenburg.pdf.
- Venter, A. J. A., & Schoonbee, H. J. (1991). The use of triploid grass carp, *Ctenopharyngodon idella* (Val.), in the control of submerged aquatic weed in the Florida Lake, Roodepoort, Transvaal. *Water SA*, 17, 321-326.
- Wells, R. D. S., & Clayton, J. S. (2005). Mechanical and chemical control of aquatic weeds: costs and benefits. Chapter 208. In: Pimentel, D. (Ed). *Encyclopaedia of Pest Management*. CRC Press 2002. <http://www.crcnetbase.com/doi/abs/10.1201/NOE0824706326.ch208>.
- Wells, R., Clayton, J., Schwarz, A. M., Hawes, I., & Davies-Colley, R. (2000). Mighty River Power aquatic weeds: issues and options. NIWA Client Report MRP00502, prepared for Mighty River Power Ltd: 55.
- Zehnsdorf, A., Hussner, A., Eismann, F., Rönicke, H., & Melzer, A. (2015). Management options of invasive *Elodea nuttallii* and *Elodea canadensis*. *Limnologia*, 51, 110–117.

Notes

1. Costs information. The assessment of the potential costs shall describe those costs quantitatively and/or qualitatively depending on what information is available. This can include case studies from across the Union or third countries.

2. Level of confidence⁸: based on the quantity, quality and level of agreement in the evidence.



- **Well established:** comprehensive meta-analysis⁹ or other synthesis or multiple independent studies that agree.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and/or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps

3. Citations and bibliography. The APA formatting style for citing references in the text and in the bibliography is used.

e.g. Peer review papers will be written as follows:

In text citation: (Author & Author, Year)

In bibliography: Author, A. A., & Author, B. B. (Publication Year). Article title. *Periodical Title*, Volume(Issue), pp.-pp.

(see <http://www.waikato.ac.nz/library/study/referencing/styles/apa>)

⁸ Assessment of confidence methodology is taken from IPBES. 2016. Guide on the production and integration of assessments from and across all scales (IPBES-4-INF-9), which is adapted from Moss and Schneider (2000).

⁹ A statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.